

U. S. PLANT PATENT APPLICATION OF

MICHAEL R. HEFFNER

FOR: MINI IMPATIENS PLANT NAMED

‘BODLIZPIN’

HEFFNER, Michael R.

TITLE: MINI IMPATIENS PLANT NAMED 'BODLIZPIN'

APPLICANT: MICHAEL R. HEFFNER

BOTANICAL CLASSIFICATION/CULTIVAR DESIGNATION:

Impatiens walleriana cultivar Bodlizpin

5

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct cultivar of Mini Impatiens plant, botanically known as *Impatiens walleriana*, and hereinafter referred to by the name 'Bodlizpin'.

10

The new Mini Impatiens is a product of a planned breeding program conducted by the Inventor in Lompoc, California. The objective of the breeding program was to develop new compact and freely-branching Mini Impatiens cultivars with numerous small flowers and attractive flower and foliage coloration.

15

The new Impatiens originated from a self-pollination made by the Inventor in December, 1999, of a proprietary *Impatiens walleriana* selection identified as code number 01IM276-1, not patented. The

HEFFNER, Michael R.

cultivar Bodlizpin was discovered and selected by the Inventor as a flowering plant within the progeny of the stated self-pollination in a controlled environment in Lompoc, California.

Asexual reproduction of the new cultivar by terminal cuttings
5 taken in Lompoc, California, has shown that the unique features of this new Impatiens are stable and reproduced true to type in successive generations of asexual reproduction.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and are
10 determined to be the unique characteristics of 'Bodlizpin'. These characteristics in combination distinguish 'Bodlizpin' as a new and distinct Mini Impatiens cultivar:

1. Compact, upright, outwardly spreading and mounded plant habit.
- 15 2. Freely branching habit; dense and bushy growth habit.
3. Small dark green-colored leaves.
4. Freely flowering habit.

5. Small light pink-colored flowers that are positioned above and beyond the foliage.

Plants of the new *Impatiens* differ primarily from plants of the parent selection in the following characteristics:

- 5 1. Plants of the new *Impatiens* are more compact than plants of the parent selection.
2. Plants of the new *Impatiens* have smaller leaves than plants of the parent selection.
3. Plants of the new *Impatiens* have smaller flowers than
10 plants of the parent selection.
4. Flowers of plants of the new *Impatiens* are light pink in color whereas flowers of plants of the parent selection are coral pink in color.

The new *Impatiens* can be compared to the *Impatiens*
15 *walleriana* cultivar Fifty Blush Pink, not patented. However, in side-by-side comparisons conducted in Lompoc, California, plants of the

HEFFNER, Michael R.

new Impatiens differed from plants of the cultivar Fify Blush Pink in the following characteristics:

1. Plants of the new Impatiens were more freely branching than plants of the cultivar Fify Blush Pink.
- 5 2. Plants of the new Impatiens had lighter green-colored leaves than plants of the cultivar Fify Blush Pink.
3. Plants of the new Impatiens had larger flowers than plants of the cultivar Fify Blush Pink.
4. Plants of the new Impatiens had lighter pink-colored
10 flowers than plants of the cultivar Fify Blush Pink.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type.

15 Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new Impatiens. The photograph at the top of the

HEFFNER, Michael R.

sheet comprises a side perspective view of a flowering plant of 'Bodlzipin'. The photograph at the bottom of the sheet is close-up view of typical flowers and leaves of 'Bodlzipin'.

DETAILED BOTANICAL DESCRIPTION

5 Plants of the cultivar Bodlzipin have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity, without, however, any variance in genotype. The
10 aforementioned photographs and following observations and measurements describe plants grown in Lompoc, California, under commercial practice in a polycarbonate-covered greenhouse during the winter and spring with day temperatures ranging from 18 to 24°C, night temperatures ranging from 16 to 18°C and light levels ranging from 4,000 to 8,000 foot-candles. Rooted young plants were planted
15 in 10-cm containers and had been growing for about 16 weeks when the photographs and the description were taken. In the following description, color references are made to the Royal Horticultural

HEFFNER, Michael R.

Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

BOTANICAL CLASSIFICATION:

- 5 *Impatiens walleriana* cultivar Bodlispin.

PARENTAGE:

Self-pollination of a proprietary *Impatiens walleriana* selection identified as code number 01IM276-1, not patented.

PROPAGATION:

- 10 Type cutting: Terminal cuttings.

Time to initiate roots: About 14 days at 22°C.

Time to produce a rooted young plant: About 28 days at 20°C.

Root description: Fine, fleshy; white in color.

Rooting habit: Freely branching.

- 15 PLANT DESCRIPTION:

General appearance: Compact, upright, outwardly spreading and mounded plant habit.

Growth and branching habit: Moderately vigorous and freely-
branching growth habit with about 12 basal branches; lateral
branches potentially forming at every node; dense and bushy
growth habit. Pinching, that is removal of the terminal apex, is
5 typically not required, but will enhance branching.

Plant height: About 10 cm.

Plant diameter or spread: About 28 cm.

Lateral branches:

Length: About 12 cm.

10 Diameter: About 5 mm.

Internode length: About 1.5 cm.

Texture: Smooth, glabrous.

Color: 146C.

Foliage description:

15 Arrangement: Alternate, simple.

Length: About 3 cm.

Width: About 1.8 cm.

Shape: Elliptical.

Apex: Acute.

Base: Attenuate.

Margin: Slightly crenate with ciliation.

5 Texture, upper and lower surfaces: Smooth, glabrous.

Venation pattern: Pinnate.

Color:

Developing foliage, upper surface: 146A.

Developing foliage, lower surface: 146B.

10 Fully expanded foliage, upper surface: 147A.

Fully expanded foliage, lower surface: 147B.

Venation, upper surface: 146B.

Venation, lower surface: 147B.

Petiole:

15 Length: About 1.5 cm.

Diameter: About 2 mm.

HEFFNER, Michael R.

Texture, upper and lower surfaces: Smooth,
glabrous.

Color: 146C.

FLOWER DESCRIPTION:

- 5 Flower type and habit: Numerous small single flowers that are
roughly triangular in shape. Four to five flowers arranged in
axillary clusters. Freely flowering, usually about 40 flowers and
flower buds per lateral branch. Flowers positioned above the
foliage and typically face upright or outward. Flowers last about
10 five to seven days under greenhouse conditions. Flowers
persistent. Flowers not fragrant.

Natural flowering season: Flowering is continuous through the
spring and summer.

Flower size:

- 15 Height: About 2.5 cm.

Width: About 2 cm.

Depth (not including spur): About 8 mm.

HEFFNER, Michael R.

Flower buds (at stage of showing color):

Length: About 8 mm.

Diameter: About 4 mm.

Shape: Ovoid.

5 Color: 69C.

Petals:

Quantity/arrangement: Five per flower in a single whorl.

Length, banner petal: About 1 cm.

Length, lateral petals: About 1 cm.

10 Length, basal petals: About 1.4 cm.

Width, banner petal: About 1.2 cm.

Width, lateral petals: About 7 mm.

Width, basal petals: About 8 mm.

Shape: Obovate to slightly cordate.

15 Apex: Rounded and slightly emarginate.

Base: Attenuate.

Margin: Entire.

HEFFNER, Michael R.

Texture, upper and lower surfaces: Smooth, glabrous.

Color:

When opening, upper surface: 75A.

When opening, lower surface: 69A.

5 Fully opened, upper surface: 75A to 75B; color becoming closer to 75C to 75D with development; towards the base, 74C.

Fully opened, lower surface: 69B.

Sepals:

10 Quantity/arrangement: Five in a single whorl; one modified into an elongated spur.

Calyx length: About 5 mm.

Calyx diameter: About 5 mm.

Shape: Elliptic.

15 Apex: Acute.

Margin: Entire.

Texture, upper and lower surfaces: Smooth, glabrous.

HEFFNER, Michael R.

Color, upper and lower surfaces: Close to 144A.

Spur length: About 2.5 cm.

Spur diameter: At flower, about 2 mm; at apex, less than 1 mm.

5 Spur color: 55D.

Peduncles:

Length: About 1.5 cm.

Diameter: About 2 mm.

Angle: About 45° from vertical.

10 Strength: Strong.

Texture: Smooth, glabrous.

Color: 144C.

Reproductive organs:

Stamens:

15 Quantity: Five fused around stigma.

Anther length: About 2 mm.

Anther color: 72D.

HEFFNER, Michael R.

Pollen amount: None observed.

Pistils:

Quantity per flower: One.

Pistil length: About 3 mm.

5 Stigma shape: Rounded.

Stigma color: 77A.

Style color: 147A.

Ovary color: 147A.

Seed/fruit: Seed and fruit production have not been observed.

10 DISEASE/PEST RESISTANCE:

Plants of the new Impatiens have not been observed to be resistant to pathogens and pests common to Impatiens.